

## Carnation Tortrix Moth Egg Killer Application Instructions

Please use immediately upon receipt for best results

## **Required Conditions:**

- Temperature range: 15-35°C
- For indoor use only
- Do not apply Trichogramma if insecticides have been used in the past 1-2 weeks

## **Before Application:**

- Do not open the sachet. The sachet already contains a small emergence hole
- Manually remove any caterpillars before placing down the sachets
- Ensure the sachets are away from any Carnation Tortrix Moth Pheromone Traps so the wasps do not become caught. The wasps are not attracted to the pheromone.

## **Application:**

- Place the sachets on a low level, horizontally, so the wasps will not be disturbed
- Ensure to avoid blocking the emergence hole
- Do not not place objects on top of sachets as you may damage the pupae inside
- One sachet can treat an area of up to 25 square metres

## **Next steps:**

- The parasitic wasps will emerge over a 2-4 week period. Replace after 4 weeks
- Observe pest levels and applying more
   Trichogramma if pest populations remain

















# Long Lasting Pheromone Trap

#### For The Monitoring & Capture Of:

#### Codling Moth / Pea Moth / Plum Moth / Leek Moth









#### Horse Chestnut Leafminer / Carnation Tortrix Moth







#### Pack Contents:

- Re-usable pheromone trap
- Species specific pheromone lure(s)



#### **Assembly Of Trap:**

- 1. Remove paper from yellow sticky pad and place at bottom of bucket
- 2. Attach the funnel to the bucket by aligning the slots on the funnel with the tabs on the bucket and twisting to lock together.
- 3. Attach the lid to the funnel by aligning the 4 inserts with the 4 pegs on the funnel. Push the pegs into the holes so the lid is fully attached.
- 4. Remove a pheromone lure from its sachet and place in the plastic cage, then fasten the cap onto the cage. Store additional lures in the fridge.
- 5. Push the lure cage through the hole in the lid from above so that the cage effectively hangs below the lid.
- 6. Thread the supplied hanging cord through the two tab-holes on the lid and tie at both ends.

#### How To Use:

- Place traps out from the end of April until the end of October
- One trap will cover over 250sqm (or one per tree if treating Horse Chestnut Leafminer)
- Hang traps at 1.5m height (or within tree canopy if treating Horse Chestnut Leafminer)
- Replace lures as per website directions and empty trap contents. Store additional lures in the fridge or freezer until required

Cap Pheromone lure Plastic cage Lid



**Funnel** 





Dragonfli Ltd, Unit 5, Cherry Tree Farm, Sible Hedingham, Essex, CO9 3LZ